P L A T E CCXCV.

ARUM, Tourn. Inst. R. H. 158. Tab. 69. Raii Meth. Plant. 74. Lin. Gen. Plant. 915. Wake-Robin, or Cuckow-Pint; in French, Pied de Veau.

The Characters of this Genus are,

THE Flower bath a large oblong Spatha (or Sheath), marked a, which is closed at the Bottom, compressed in the Middle, and open at the Top, as is represented at b: The Spadix (or Stalk), which sustains the Leaves and Flowers, is single, marked n; upon the Top of which are situated the Male Flowers, as is shewn at c; and below these are the Germina, marked d. These Flowers have no Petals, or Stamina. One of the Male Flowers is represented of the natural Size at e, which is magnified at f: At g is shewn an Embryo of its natural Bigness, and one which is magnified at h. The Embryo with its Cover is represented of its natural Size at i, and another magnified at k; and at 1 is shewn the naked Embryo of its natural Size; and at m is another magnified. These Germina afterward become Berries, with One Cell, inclosing a single Seed.

This Genus of Plants is ranged by Tournefort in his Third Class of Plants, which includes the Herbs with an anomalous Flower of One Leaf, shaped like an Ear or Hood. Mr. Ray places it in his Seventeenth Class, which includes the Herbs bearing Berries. Dr. Linnaus ranges it in the Seventh Section of his Twentieth Class, which includes those Plants whose Flowers have many Stamina, which sit upon the Style or Pointal.

The Specie here represented is,

Arum caulescens, foliis sagittatis, Lin. Sp. Plant. 967. Stalky Arum, with arrow-pointed Leaves. In the Gardener's Dictionary it is titled, Arum caulescens, foliis sagittatis, spatha declinata clausa; Stalky Arum, with arrow-pointed Leaves, and a declining closed Sheath. Plumier titles it Arum arborescens, sagittariæ soliis. Plant. Amer. 44. Tree-like Arum, with arrow-pointed Leaves: And Sir Hans Sloane, in his History of samaica, calls it Arum caule geniculato, cannæ Indicæ soliis. Cat. 63. Arum with a jointed Stalk, and Leaves like the Indian Cane. In the West-Indies it is commonly called Dumb Cane.

This Plant rifes, with a jointed green Stalk, to the Height of Eight or Ten Feet, and is about the Size of a large Waskin z-frick: The Leaves come out toward the Top, in an irregular Order; they are oblong, of a lightgreen Colour, and are from Seven to Ten or Eleven Inches long, and Five or Six broad in the Middle, having Two Ears at their Base, like the Beards of an Arrow. Their Footstalks embrace each other at their Base, and between these come out the Spathæ, or Sheaths, as is represented at a. They are Five or Six Inches long, of the same Colour with the Leaves, swelling at a small Distance from the Footstalk, and compressed again in the Middle, opening a little at the Top, where the Point of the Spadia appears, upon which the Flowers are irregularly placed: The fe at the Top are formed into a close Spike, but those below are ranged sparfedly. The Sheath is closely uxed on one Side to the Spadix, fo cannot be separated from a without being forn; but on the opposite Side it may be open to so as to snew the Spadix, with its Embryo, as is represented of d; and it is only upon this Side of the Space that the lowers are ranged, for the Sheath Continuers to the other. The Embryo on the lower Part of the Spadix have each a fmall Cover, compoied or Five very fmall white Leaves, which are fafte ed to the under Part, and separate above, as is shewn at i and k. The Embrya are green, kidney-shaped, and have brown Covers. The Male Flowers, at the Top of the Spadix, are of an irregular square Figure, and are brown. The Embrya do not come to Maturity in England; the utmost Perfection they have yet arrived to is as that represented in its natural Bigness at I, and is magnified at m. The Time of its Flowering is in the Spring of the Year; the whole Plant is filled with a sharp acrimonious Juice, which will blister the tender Skin, if applied to it; and, if a Part of the Leaf is broken and applied to the Tip of the Tongue, it occasions a very painful Sensation, causing the Salivary Ducts to swell, and brings on a great Defluxion of Saliva: The Inhabitants of the Sugar Islands do sometimes rub the Mouths of their Negroes with the Juice of this Plant, by way of Punishment; and as this swells their Mouths, so as to prevent their eating or speaking till the Swelling is asswaged, so it is called by them Dumb Cane.

P L A T E CCXCVI.

Dracontium, Lin. Gen. Plant. 916. Arum Plum, Amer. 40. Tab. 56. Dracunculus, Tourn. Inft. 160. Tab. 70. Dragon; in French, Serpentaire.

The Characters of this Genus are,

once, with One Valve, represented at a, and a single cylindrical Spadix, on the upper Part of which the Parts of Interification are disposed in a singular manner, each being closely sastened to the Spadix, as is shewn at b, having no Empalement, nor any Petals which can be properly so called. In this Specie there are but Iwo Stamina in each Flower, which have spatte-shaped Summits, represented at c and d of their natural size, and at c c magnified. I he Germen c is shaped like Number 14.

a Vase, baving a coloured Crown, which is shown magnified at A e, and at A there are Three of the Germina cut transversely, wherein are seen the Embrya of Two Sceds.

The Specie here represented is,

DRACONTIUM foliis pertufis, caule scandente, Lin. Sp. Plant. 968. Dragon with perforated Leaves and climbing Stalks. Piumier titles this Plant, Arum bederaceum, amplis foliis perforatis. Piant. Amer. 40. Tab. 56, 57. Climbing Cuckow-Pint, with large perforated Leaves.

This Plant grows naturally in many of the Islands in the II est Indies, where the Stalks fasten themselves to E e e

the Trees which are near them, by their Roots, which are put out at each Joint, and fix closely to the Bark of the Trees; so that, if the Stalks are cut off below, these Roots will draw Nourishment sufficient for the Growth of the Plant, from the Trees to which they are fastened; so, by this means, the Plants will rife to the Top of the highest Trees. In Europe, where they must be kept in a warm Stove, if the Plants are placed near the Wall, the Roots which come out from the Stalks will fasten to the Wall, and thereby mount to the Top of the Stove; but those Roots which come out on the opposite Side to to the Wall, will hang down several Fect; and if, in their Way downward, they meet with any Pot of Earth, they will fasten themselves into it, or to any other Support near them. The lower Part of the Stalks are feldom larger than a Goose-quill; but that Part of the Stalk which immediately sustains the Flower, is often as thick as a Man's Thumb. They are cylindrical, of a deep-green Colour, jointed, having a fingle Leaf at each Joint, whose Footstalks do almost embrace the Stalks with their Base. The Leaves are from Five to Seven or Eight Inches long, and from Three to Four or Five broad. On each Side of the Midrib are Two or

Three Perforations of an oblong Form, which appear as if they were eaten by some Intects, as is represented at g. The Spadix arises at the Top of the Stalk, and is embraced by the Base of the Footstalks of the Leaf: This has a large boat-shaped Spatha (or Sheath), which opens on one Side the whole Length; and when the Parts of Fructification arrive near to Maturity, the Spatha (or Sheath) separates from the Spadix, leaving it uncovered in the manner as is here represented, having Marks of the Parts of Fructification on the Infide, which appear like to many Spots. The Time of this Plant's Flowering is not constant; sometimes they appear in the Spring, and at others in the Autumn; but they are of long Duration, frequently continuing Five or Six Months: But the Fruit does not come to Maturity in England.

As there has not been any good Figures of these Two Plants, wherein the Parts of the Fructification have been well represented; and these having, for some Years past, produced their Flowers in the Chelsea Garden, so I have taken the Liberty to exhibit the Plants in the

best State to which they have arrived here.

PLATE CCXCVII.

Fig. 1. Gladiolus. Fourn. Rey. Lin. &c. Cornflag; in French, Geicul.

A S the Characters of this Genus of Plants has been already exhibited in the Course of this Work, it is unnecessary to repeat them here.

The Specie here represented is,

GLADIOLUS, foliis ensiformibus, floribus alternis, corollis patentibus. Cornflag, with sword-shaped Leaves, Flowers ranged alternately, and spreading Petals.

This Plant grows naturally at the Cape of Good Hope, from whence the Seeds were fent to Holland, where the Plants were raifed by Mr. Willem Vanhazen, near Leyden, who was so kind as to send me a Root of it.

The Root is tuberous and compressed, having a greyish Skin or Cover, from which arise Four or Five narrow fword-shaped Leaves, Nine or Ten Inches long, and about One Inch broad in the Middle, where they are broadest, as is shewn at a; of a deep Green, ending in acute Points. Between the Leaves come out the Flower Stem, which is cylindrical, and rifes a Foot and a Half high, represented at b, garnished with Two or Three fmaller Leaves, according to the Number of Joints in the Stem, which is almost embraced by their Base. The Flowers are ranged alternately toward the Top of the Stalk, each being inclosed in a Spatha (or Sheath) of One Leaf, which opens on one Side, as is represented at c: The Flower is of One Leaf, which is deeply divided into Six unequal Parts, as is represented at the Bottom, and has Three awl-shaped Stamina, which are interted at their Bafe, alternately, into every other Segment of the Petal, represented at a. The Germen, which is fnewn arg, is fituated under the Flower, supporting a flender Style, which is longer than the Stamina, crowned by a triffed Stigma, represented at f. The Flower is of a pale Flesh Colour, and appears in May. After the Flower is paft, the Germen swells to an oblong blunt Capfule, having Three Cells, which are filled with irregular shaped Seeds. Soon after the Seeds are ripe, the Stalk and Leaves decay, and the Roots remain

inactive till Winter, when they put out new Fibres, and the Leaves push out early the following Spring.

Fig. 2. WATSONIA.


The Characters of this Genus of Plants have been exhibited already in the Course of this Work, so need not to have been repeated here, were it not for the Alteration of the Title to this Genus. Dr. Trew, of Nuremburg, who had a Drawing of the First Sort before represented in this Work, sent him, which was taken from a Plant in Flower in the Chelsea Garden, published it among his rare Plants, under the Title of Meriana, before he had seen my Book, in which I had given it the Title of Watsonia, in Honour of Dr. Watson, F. R. S. who is a very learned Botanist: But upon seeing my Title, he wrote to me, that he would suppress his in his Tables, and continue mine to the Plant, as I had raised it from Seeds, and had it first in Flower in the Chelsea Garden: But since this, Dr. Linnaus, in his last Edition of the Systema Natura, has ranged it under the Genus of Antholyza, to which he has also added the Cursuia of Butner. But as both these Plants do essentially differ in their Characters from that which he has chosen to constitute his Genus, the Flowers of which have but One Stamen, and both these have Three; so there is Reason for keeping them separate: And as the Flowers of Cunonia are divided almost to the Bottom, and are shaped like a Buttersly, so that should not be joined with this Plant, whose Flowers have long Tubes, the Brims of which are cut into Six equal Parts, fpreading open. This Shape of the Flower also removes it from the Gladiolus, to which it is nearer allied than to the Antholyza; therefore I shall continue my former Title to the Genus, and believe those who will examine the Characters with Impartiality, will agree with me to to do.

The Specie here represented is,

Warsonia bunilis, foliis lineari-ensiformibus, tubo floris longissimo. Dwarf Watsonia, with linear sword-shaped Leaves, and a very long Tube to the Flower.



Plug I Gravia State ensignmilias floribus atternis, corollis valentibus, 2 Plug 2 W VVSOXIX Tunnitio, polis lineari ensignmeilino, tako floris longigimo.



TRILMFETTA plus dilengo cratos elituje ferralis petrello lengificimo.

I Miller be et acuip

". i. let according to det by " tiller july sole; to . -



SYDEROXYLUM min. mem.

M. Willer det & de

Dublishil according to the elet, by &Poulle in oso, 1760. -

This Plant is a Native of the Cape of Good Hope, from whence the Seeds were brought by Captain Hutchinson, of the Godolphin Indiaman, in the Year 1754. The Root is tuberous, compressed, and covered with a brownish Skin; from which come out Four or Five narrow sword-shaped Leaves, about Five or Six Inches long, and One-third of an Inch broad in the Middle, ending in acute Points, of a dark Green, embracing each other with their Base, see Letter a. From between these comes out the Flower Stem b, which is naked, and rises a Foot high: Toward the Top the Flowers are ranged alternately, each having a Spatha (or Sheath), which is divided into Two Parts, as is represented at c. The

Tube of the Flower is long, narrowed at the Base, but is greatly enlarged above, as is shewn at d. The Brim is cut into Six equal Segments, as is represented at e, which spread open. Under the Flower is situated the Germen, supporting a slender Style, crowned by a trisid Stigma, each Division being bisid and reslexed, as is shewn at g. This Style is attended by Three slender Stamina, of the same Length, represented f f, terminated by moving Summits. The Flower is of a bright Red on the Outside, but is paler on the Inside. These appear the latter End of May, and continue in Beauty a Fortnight or Three Weeks; then the Leaves and Stalk decay, without producing Seeds here.

PLATE CCXCVIII.

TRIUMFETTA, Plum. Gen. 40. Tab. 8. Lin. Gen. Plant. 529. Bartramia Lin. Gen. Plant. 480. We have no English Title for this Plant.

The Characters of the Genus are,

THE Empalement of the Flower is cut into Five linear Segments a, which foon fall off; it has, for the most Part, Five narrow Petals b, which are erect, and Ten slender Stamina c, which are terminated by roundish Summits d: In the Center of the Flower is situated a roundish Germen, supporting a hairy Style c, the Length of the Stamina, crowned by a single Stigma: The Germen afterward becomes a roundish prickly Fruit, whose Spines are crooked f, having Four or Five Cells g, each containing One Seed h.

The Specie here represented is,

TRIUMFETTA, foliis oblongo-ovatis, obtuse serratis, petiolis longissimis. Triumfetta, with oval oblong Leaves, bluntly sawed, standing upon very long Footstalks.

This Plant grows naturally in *India*, from whence the Seeds were brought, which have fucceeded in the *Chelfea* Garden, where the Plants have flowered and perfected their Seeds. It is annual, and perifhes foon after

the Seeds are ripe, so is very different from that Specie which Doctor Linneus titled Bertramia, in the former Edition of his Genera; but in the last Edition of his System he has joined it again to Triumsetta, from which it only differs in having sometimes Ten Stamina, whereas the Triumsetta of Plumier has rarely more than Eight. When the Flowers of this Plant has but Four Petals (or Segments), then there are but Eight Stamina in the Flower; but when the Flower has Five Petals, then there are always Ten Stamina in each.

The Plant rifes about Two Foot and a Half high, fending out several Branches on every Side, which are garnished with oval Leaves, ending in long acute Points, some of which are heart-shaped, and others have an Angle on each Side towards the Point. They are from Three to Four Inches long, and almost as much in Breadth in their broadest Part, standing upon very long Footstalks, and are crenated on their Edges.

The Flowers come out in long loose Spikes at the Top of the Plant, being ranged without Order. They are small and yellow, some having Four Petals, and Eight Stamina, and others have Five Petals and Ten Stamina. After the Flower is past, the Germen turns to a roundish burry Fruit, with Four Cells, each containing One Seed. It slowers in September, and the Seeds ripen in Nevember.

PLATE CCXCIX.

Siding oxygum. Dillen. Hort. Edib. 263. Lin. Gen. Plant. 234. Iron-wood.

The Characters of the Genus are,

It but Male and Hermaphrodite Flowers on different Plants. The Male Flowers have an Empalement, which is cut into Five obtage Segments with the Flower is divided into Five roundiffs concave Parts V., almost to the Bottom: It has Tive incurved Stamina A, which are terminated by oval Summits v. In the Center is placed a roundiffs com-

pressed pointed Body, resembling a Germen, I, which has neither Style or Stigma, and falls away with the Flower, without having any Appearance of Fruit.

This Genus is placed in the First Section of Linneus's Fifth Class, which includes the Plants whose Flowers have Five Stamina and One Style: But as the Male Flowers grow on separate Plants from the Fruit, so it should be referred to the Twenty-third Class. But as Doctor Linneus has not seen the Flowers of these Plants, so he may be excused for ranging them here. The thorny Sort, which is common in the West Indies, also agrees with this, having Plants which bear only Male Flowers, Two of which have produced Flowers in the Chelsea Garden; and Dr. Houstown sent Specimens of the Male and Hermaphrodite Trees from Jameica to the Author.

The Sort here represented is,

Sideroxylum mas inerme. Male Iron-wood without Thorns.

This Tree grows naturally at the Cape of Good Hope, where it rifes to the Height of our Apple-trees in England. The Wood of it is so heavy as to fink in Water. But in Europe the Trees do rarely arrive to more than Eight or Ten Feet high, dividing into many Branches, which are covered with a Russet Bark, and are garnished with spear-shaped Leaves, about Three Inches long, and an Inch and a Half broad in the Middle, ending in Points at both Extremities. They are placed without Order on the Branches, having Footstalks an Inch long. The Leaves are smooth, and of a lucid Green, continuing all the Year. The Flowers come out in Clusters from the Side of the Branches, standing upon short Footstalks, which branch out into several smaller, each suftaining a single Flower, which is small and white, one of which is represented of its natural Size at g, the other is magnified. This Plant slowers in May and June.

PLATE CCC.

Lycium, Lin. Gen. Plant. 232. Jasminoides Nissel. Ac. Reg. Par. 1711. Rhamnus C. B. P. 477. Boxthorn.

HIS Genus of Plants is ranged in the First Section of Linneus Fifth Class, which includes those Plants whose Flowers have Five Stamina and One Style. As the Characters of this Genus have been already exhibited in the Course of this Work, so it is needless to repeat them here.

The Specie here represented is,

Lycium, foliis cordato-ovatis, oppositis sessilibus perennantibus spinis crassis bigeminis, floribus confertis. Dist. Hort. Boxthorn, with oval heart-shaped ever-green Leaves, placed opposite, sitting close to the Branches, thick Spines in Pairs, and Flowers in Clusters.

The Seeds of this Plant were brought from the Cape of Good Hope to the Dutch Gardens, where the Plants were first raised, and have since been communicated to the curious Botanic Gardens in most Parts of Europe. Some of the Dutch Botanists have ranged this Plant under the Genus of Randia, from the Shape of the Flower, to which it has a near Assinity, if the I ruit is not admitted as a Character; but, if this be allowed, it must be ranged with the Lycia, for the Fruit of the Randia has but One Cell, which is filled with cartilaginous compressed Seeds; whereas the Fruit of this has Two Cells, each containing a single Seed. It is a low Shrub, which seldom rises above Four or Five Feet

high, sending out Branches from the Root upward, which divide in Pairs and spread asunder. These are covered with a green Bark, as is shewn in the Figure, and are garnished with oval heart-shaped stiff Leaves, about an Inch long, and Three Quarters of an Inch broad at their Base, ending in acute Points, sitting close to the Branches, and are placed opposite; they continue green all the Year. At the Divisions of the Branches stand the Spines, which are forked and double: They are short and stiff, for the most Part growing erect. At the End of the Branches the Flowers are produced in Clusters, as is represented at n. The Cup of the Flower, shewn at a, is small, erect, and slightly indented at the Top in Five Parts; this is permanent. The Flower is funnel-shaped; the Tube b is cylindrical, and a little incurved at the Top, where it is cut into Five acute Segments, which spread open, represented at c. In the Tube of the Flower there are Five awl-shaped Stamina, terminated by erect Summits, as at d. The Germen is situated in the Flower Cup, supporting a single Style, which is longer than the Stamina, shewn at e, and crowned by a thick bifid Stigma, marked f. After the Flower is past, the Germen swells to an oval fleshy Berry, represented at g, which has Two Cells, as is shewn at b, where the Berry is cut transversely at i, in each of which is lodged a fingle oblong hard Seed, reprefented at k.

This Shrub produces its Flowers in July and August, which are of a clear White; and the Fruit ripens the Spring following, which is of a deep-red Colour when ripe.



LAVAV M. jetins cordate courtes oppositis fefritibus perennanutius, opines crajais vigeminis gioritius conjectis. Diet Hort.—

L. Mitter det 8 se.

Publishit secontem with it to be 2. Mitter fine 80% gio.—

INDEX PLANTARUM

Quæ in hoc opere exhibentur.

A		Fig.	Pag.	1			Fig.	Pag.	•		Fig.	Pag.
A BIES -		1	1		В		- 0		Cucubalus	-	112	75
Abrotanum .		2	2	Balfamina	-	-	59	•	Cunonia		11.3	75
Abutilon —	-	3	3	Basteria Bauhinia		-	60		Cyanus — Cyclamen —	-	114	76
Acacia non spinosa — Americana	_	- 4	4	Belladona			61 62	41	Cyclatilei	-	116	7 7 7 7
tenui folia		6	4	Berberis	terrang .		63	42	Cypripedium	-	242	161
foliis bispinnatis		291	194	Bidens	-		64	43	Cytifus —		117	78
Acanthus ——		7	5	Bignonia	-		55	43				
Acer —		8	5	Bistoria			67	44	D'ayena —		T T Q	70
Achillea		9 10	7	Blattaria Borrago			_	45	Dayena		5119	79 79
Achyranthes -		11	7	Browallia	-		69	46	Delphinium -		250	167
Aconitum —	•	12	8	Brunella	(0.0000)		70	47	Dianthus -		\$121	} 81
Adhatoda		13	9	Bryonia			71	47			L122	١ .
Adonis		14	9	Buglossum Bulbocodium			7 ² 240		Dictamnus —		123	82 83
Agave ————————————————————————————————————	_	222	10			(73		Digitalis	_	120	80
Alaternus	-	16	11	Bupthalmum		1	249	166	Diosima	-	125	83
Alcea	_	17	II	Bupleurum	-	-	74	49	Diospyros	-	126	84
Alchimilia —		18	12				•		Dodartia ——	_	127	85
Aloe	. {	19	13		. C	c	he	-	Doronicum —— Dracocephalum		128	85
Alysfon —	_	292	195	Calendula		<u> </u>	75 76	50 51	Dracontium -		296	197
Amaranthoides —	-	21	14			. 1		189				
Amaranthus —		22		Cannabis	_		77	51	. E			
Amaryllis	્ક	23		Capnoides		-	78		Echinops		130	87
	ŧ	24		Caprifolium Cardiaca			79		Elichrysum —		131	87 88
Ammi — Amomum —	_	25 26		Carduus				53	Epimodium —	******	133	89
Amorpha —	-	27		Caryophyllus			81	54	,		55	
Amygdalus —	-	28		Cassia			82	55	F			_
Anchula —	-	29		Cassine		-	83		Fabago	-	134	89
Androface —		30		Castanea Caucalis			84 85	_	Ferraria —— Filago ——		280	187
Anemone — Anguina —	_	32		Ceanothus			86	57 57	Fragaria		135 288	192
Anguria —		33		Celastrus			87	58	Fumaria	-	136	91
Anil —		34		Celtis		-	88	59				
Anona —		35	_	Cerafus		-	89	59	Cologa	•	10~	^-
Anonis —		36		Cereus Cerinthe		_	90		Galega ——— Gallium ———		137	91 93
Anonis viscosa — Anthemis —		37 28	25	Chelidonium	-		92		Genista ——		138	92
Anthericum —		39		Chelone		-	93	62	Gentiana .	-	139	93
Antholyza		40		Chironia			97	65	Geranium —	_	140	94
Anthyllis —		41		Cirfium					Geum -		141	94
Antirrhinum — — Aphaca —		42		Clematis Clethra			284 281	. 2 -			235	9 5
Apocynum —		43 44	29	Clinopodium	-	-	95	63	Gladiolus		236	158
Apocynum ereclum	*******	45	30	Clymenum	-		96	64			297	198
Aquifolium	_	46	31	Cncorum	dere-ord	-	98	65	Glaucium —	-	143	96
A quilegia —		47		Cnicus	(Martine Park)			66	Gundelia	لنستسق	287	191
Arctoris —		48 49	33	Colutea	-	- 3	99	67	H			
Argemone	-	40	33	Convallaria				67	Helianthemum	-	290	193
Aristolochia		51	34	Convolvulus	-		102		Heliotropium	-	144	96
Arum		52	34	Conyza	_	-			F-Ielleborine —		145	97
	-	295	197	Corona Impe	rialia		104		Hieracium majus		149	97 98
Afarina I farum		53	35	Corona Impe	f heart's	-	106		Hippocrepis —		278	185
A (clepias		53	35	Coronilla		{	107	71	I-lyacinthus		148	98
Afcyrum	****	54	36					193	Hydrangea		251	167
Afparagus	-	55	~ Q	Craffula					Hypecoon	-	149	100
Atphodelus —— Atter Carolinianus	patient)	50	38 38	Cratægus	-		109 269	13 170	Hypecoon — Hypericum —	-	150	101
Atter, caule corymbolo		57 282		Crinum				73				
Aftrigalus		58	39	Crocus	,		111	7+	Fff		I.	Jacea

INDEX PLANTARUM.

I	Fig. Pag. 101		£ 188	Pag.	Sarracena —	Fig. 241	161
Jacea ———	- 1153 102	Ochothera —	189	126	Saxifraga	243	
Jasminum	- 180 120	Omphalodes —	- 190	127	Schinus	- 240	_
Iris ——	- 154 103	Opuntia —	- 191	127	Scolymus —	- 229	
		Ornithogalum	 192	128	Scorpiurus	- 230 - 237	
Ixia ——	-< 156 10+	Orobus		129	Scrophularia —	- 23I	_
	237 158	Osteospermum —	- 194	129	Securidaca —	$\frac{-}{-}$ 232	
K	_	Othonna	-		Senecio ——	5234	
Kalmia ——	<u> 228 152</u>	Othonna —	 245	163	Serratula	一 { 247	
		Oxalis	- 195	130	Siliquastrum	- 253	
L		D			Sideroxylum —	- 299	-
Lactuca	- 157 105	73 1	- 106	131	Solanum	- 294	
Lamium —	- 158 105		— 196 — 199			§ 254	
Lappa -		Pæonia ——	— 199 — 197	132	Solidago —	- 1 255	_
Lathyrus		Pancratium —	- 198	122	Spartium	259	
Lavatera		Pavia —— Pentapetes —	- 200		Spiræa	_ {256	3 171
Leonurus		Perficaria	- 201	134	Spiræa	257	5-1-
Ligustrum —	C 160 7		202		Stapelia ——	- 258	172
Lilac	$$ $\left\{\begin{array}{c} 103 \\ 164 \end{array}\right\}$ 109	Phlomis	- 203	135	Styrax	- 260	173
Lilium	— 165 x10		204	130			
Linaria —	— 166 III	2011	205	137	T		
Linum		Physalis	- 206	127	Tacamahaca	261	174
Lonicera	<u> </u>	Phytolacca -	*	138	Tamarix -	- 262	
Lotus	<u> </u>	Pinus	- 208	139	Tetragonia -	— 263	-
Lunaria	- 169 113	Polemonium	- 209	140	Teucrium —	— 264	•
Lupinus	- 170 113	Polyanthus	210	140	Thalictrum —	205	
Lychnis	170 114	Ptelea	211	141	Tordylium	<u> </u>	
Lycium -		Pulmonaria -	- 212	141	Trifolium —	— ∠67	
Lycium —	- 300 200	Punica	- 213	142	Triumfetta -	298 268	
•					Turnera —	200	-/3
\mathbf{M}		Q "	274	* 40	v		
Magnolia —		Quamoclit —	- 214	143		S 271	3 181
Malpighia —		Quercus	- 215	-43	Veratrum	- { 272	188
Mandragora —	— 173 115 — 286 191				· ·	_ \$273	
Martynia —		Ranunculus	- 216	144	Verbascum	277	•
Meadia —	, .	Refeda	- 217	145	Viburnum	- 274	
Medicago —	- 175 II5	Rheum	- 218	145	Vinca	— 186	
Mesembryanthemum		Rhus —	- 270	180	Vitex	- 275	183
		1	5219	146			
Mespilus	C ' > 110	Ricinus	7 220	147	W		
Mimofa		Robinia	- 244	162	Warneria -	 285	192
Monarda ——		Rofa	- 221	147	Wationia	- 276	184
	[220]	Rubus	- 223	149	Wationia	— 297	198
Morea	$-\left\{ \frac{239}{240} \right\}$ 159	Rudbeckia	- 224	149			
Myrtus	- 184 123				X		-00
2.2,		S			Xeranthemum —	279	186
N		Salvia	- 225	150			
Nicotiana	<u> 185 124</u>	Sambucus	226	151			
Nigella	- 187 125	Santolina	227	151			
		1					

ENGLISH NAMES of the PLANTS

Engraved and Described in this Work.

A	Fig.	Pag.			Fig.	Pag.	Fig. Pag	c t
Cacia, without The	orns 4	4	Cassioberry	-	83	55		5. 7
Acacia, the Ameri	can s	4	Catchfly	-	170	114	PJETON PLOWER	16
Acacia, the Narro	W-16		Caterpillar	-	230			0
leaved	- ໂ o	4	Celandine —		92	61		
Acacia, the Falle -	244	163	Centaury	-	97	65	F	
Adonis Flower —	14	9	Chaste Tree -	-	275			3
Agave —	222		Cheese-Runnet	-	139		Fennel Flower 187 12	400
Agrimony — —	15		Chelone —		93		Ferraria — 280 18	7
Agrimony, the Waterhem	•	43	Cherry -		89		17:	3
Alkanet — —	29	29	Chefnut —	-	84	50	Figwort — — 231 15	+
Almond Tree	28		Chickling Pea Chickweed, Berry-be	arine	160	107	Fir, or Spruce Tree - I	ı
Aloe, African	3 19	13	Chrysanthemum H	army) 112	75	771 - 1	
Amaranthus, spiked	292	195	feeded		194	129	171 - 1 - T	9
Amaryllis — —	24	16	Climber		284	186	Clause Canala	3
	-		Clethra —					30
Anemony — — —	31		Cnicus —	-	248	165	roxgiove — 3	
Arimart —	49 201				5 1067			35
Arum	295	197	Colutea, joint-podde	d	107	7 1 1	F	; 2
A farabaca common —	53	35	Colutea Shrubby, joi	nted-)		Fumicory - 776	I
Asparagus	55	37	podded	-	289	193	E	4
Afphodel	56	38	Columbine		47	31		•
		•	Convolvulus, Scarlet	-	214	143	G	
\mathbf{B}			Corn Flag, Strange		40	27	Gentian — 139 9	13
Balsamine, Female -	59	40	Corn Flag, Greater	By-	142	0-	Germander - 264 17	-
Barbadoes Cherry —	181	121	zantine -		51+2	95		,
Barberry Bush	63	42	Corn Flag African	-	235	×57	[122]	18
Barrenwort	133	89	Corn Flag African	-	236			4
Basteria	60	▼	Corn Flag -		297			7
Bastard Lychnis	215		Cotton Weed	-	135			E
Bastard Asarum	252		Cowssip Chickweed		30	20	Golden Rod \{ \frac{254}{256} \]	19
Baulm, the Turkey —	129		Cranesbill	-	140	94	L 455 1/	O.
Bean Caper	134		Crowfoot -		216		Goldylocks — 227 15	2
Bear's Breech	7	_	Crown Imperial	-	105		Groundsel — 233 15	_
Bear's Ear —	174		Cuckow-Pint		52		Gundelia — 287 19	I
Bindweed	102		Custonia, the Scarlet		113	75	7.7	
Bird Cherry —	196	_	Custard Apple		35	23	Hare's Ear — 74 4	
Bird's Eye ———————————————————————————————————	167	9	D				I Tourse	9
Birthwort —		112	Daffodil Sea -		107	122	I Touris a superior	•
Bishop's Weed	25	34	Daffodil Lily	-	197		f.T. and and and	
Bitter Vetch	193	129	mea o		118		TT 1 1 C	8
Bladder Pea	41	4	Date Plum Indian		126		Hawthorn — 179 11	
	\$ 99	,	Dead Nettle -		158		11 1	6
Bladder Sena	(100		Dianthus	,	121	81	Litaliahana sha Da Gamil	7
Blue Bottle	114	76	Diantinus		122	8 I	17.11.1. S2717 - 0	
Borage	68	45	Diervilla		124	031	272 5	L
Boxthorn	171		Diofma	-	125	83	Hemp — 77 5	I
Boxthorn	300	200	Dittany, the White	-	123			9
Briony white	71		Dodartia		127			I
Broom	138		Dogsbane		44			3
Broom, the Green	259		Dogsbane, upright	-	45	6.0	Honeyfückle, the Upright 167 rt	2
Browallia	69		Dogwood -	-	101		Honesty — 109 11	
Buglofs — —	72		Dragon	-	296		Honeywort - 91 6	_
Bulbocodium — —	2.40		Dragon's Flead		159		Horse Chesnut, the Scarlet 198 13	
Burdock —	259	100	Dropwort		250		Hyacinth Eastern — 148 9 Hyacinth, the Tuberose 210 14	
			Dwarf Ciftus	-	290	193		
Calve's Snout	4.0	28	17				lypecoon — 251 16	,
Cammock — —	42	24	***		61	4 1	-71,0000	~
Campion		44 111	Elder		_	141	I	
Campion ————————————————————————————————————	121	87	Elder — Emony —	-	21	21	Jasmine 180 12	O
	- 3 -	- /			3		Jaimir	

ENGLISH NAMES OF PLANTS.

		-	T	1		***		4		***	
Jasmine, the Persian		Fig.	Pag.	124 11 1 2 2 1		Fig.		Spiderwort -		Fig.	
Janinine, the remain	. —	169	7	Myrtle -		184	45 123	} *		· 39 €256	20
Jerusalem Sage		203	L 135				5	Spiræa	-	257	171
		204	1 1 2 0	,	N			Spotted Sanicle	-	141	94
Jersey Tea		86	~ ~ /			88	59	Staff Tree -	_	§ 16	11
Immortal Flower		279	186	Nightshade		29+	196			2 87	
Indian Fig —— Indigo Baffard ——		191	127	Nightshade, the I None so pretty	Jeadly	62	41	Star of Bethlehem Starwort Carolina		192	128
Indigo ———		² 7 34	22			141	94	Starwort American	_	282	38 188
St. John's Wort		151	101		0		_	Storax Tree		260	173
Ironwood	-	299	199	Oak		215	143	Strawberry		288	
Judas Tree	-	253		Oil-seed —		. 219	146	Strawberry Tree		48	32
•		155		Orpine, the Lesser		108		Sumach, the Venet			180
Ixia	-	150	_	Othonna		194	_	Sun Flower, the Du	art —		149
K		L237	158	Ox-eye —		3 73	49 166	Swallow-wort		53	35
Kingspear —	-	56	38			2 249	100	Sweetwilliam		214	172
Knapweed -	-	152	101	1	P						"TJ
Knapweed -		153	102	Parsley, the Bastaro	i	85	57	7	7		
				Pellitory of Spain		38	25	Tacamahac Tree	-	261	. 174
L. Tadias Dadibasas				Pentapetes —	-	200	133	Tamarifk —		262	175
Ladies Bedstraw Ladies Slipper	-	139	93	Perriwinkle — St. Peter's wort		186		Tame Poison		53	35
Larkspur, Great Bee		242				54 14	30	Tetragonia — Thistle, the Torch		203	175
Larkspur taller, pere					*****			Thistle, Melanchol		94	
Lavatera -		-				207	138	Thistle, the Golden	-	229	-
Lavender Cotton			_		-	208	139	Thistle, Woolly		293	195
Laurel, the Rose, or M	oun-	228	152	Pink		81	54	Thorn, the Egyptia	n · —	5 4:	. 4
fulli		3	0 -	Piony —		199	133	T10		1291	194
Leopards-bane Lettuce —				Pipe Tree — Pipperish Bush		103	109	Toad-flax —		166	III
Lilac		157	100	Poke, or Pork Phys	ic —	207	128	Trefoil, the Bean		105	124
Lily		169	110	Polyanthes -		210	140	Trefoil, Carolina Sh	rub .	211	141
Lily Asphodel		109		Pomegranate -			142	Trefoil, Star-headed	i	267	178
Lily Daffodil	Street-o	23	15	Poppy, the Prickly	-	50	33	Triumfetta	-	298	199
Lily, the Mexican	-	24	16	Poppy, the Horned	l —	143	96	Trumpet Flower	•	65	43
Lion's Tail London Pride		162	102	Primrose Tree		188		Tulip Tree, the L	aurel-	172	115
Lungwort		212	2 .	Privet		162	120	leaved — Turnera —) <u>(</u>	_
Lupine —		170		Punica -		213	1	Turnsole	_	268 144	179 96
		,				3		Tyger Flower		280	187
M				Q	_		1	•			
Madwort -	_	20	13	Quince —	-	116	77	v	1	:	
Magnolia — Malabar Nut	-	172	115	D				Valerian, the Greek		209	140
Mallow, the Yellow M	[arfh	13	9	Ranunculus K		216		Venus Navel-wort Vetch Kidney		. 190	127
Mailow, the Vervain		17	~ (Ragwort foreign		245		Vetch, the Chickling		41 96	27 64
Malpighia —	-	181		Raspberry -	-	223	149	Vetch, the Horse-sho	be	278	185
Mandrake —	-	173		Restharrow —		36		Vetchling —		43	29
Mantle, Ladies	-	18	1		{	37	25				
Maple Tree		. 8				218	145	W	•		
Marigold —	}	75	~ ,	Rocket, the Bastard Rock-Rose —				Wake-Robin —		42.	34
5,14,15	1	283		Rose -				Warneria		285	190
Marigold, the Fig	_ `	176	117				_	Wationia		35 276	23 184
Martynia —		286	191	S			1	Wationia, Dwarf		297	198
Mastick Tree, Indian	_	246	2.1	Saffron —	-	III		Water Lily -		239	159
Maudlin —	{	9	6	Sage		225	150	Wayfaring Tree	******	274	183
Meadia		10		Sawwort		234	156	Whin, Petty -		36	24
Meadow Rue	-	265	110	Saxifrage -			162	White Satin —— Widow-wail ——		169	113
Meadow-fweet		257	171	Self-heal	(Prophose)		1		(98	65
Medlar —				Senna, the wild -		82	55	Wild Service		269	73
Melon, the Water		33		Senna, the Bladder	-	100	67 1	Vinter Cherry	_	206	137
Melembryanthemum	-	177		Senna, the Scorpion		132		Wollsbane —	-	12	8
Milk-vetch	-	58		Senfitive Plant	-		. 1	Wood-forrel		195	130
Monk's Hood		113	1	Side-faddle Flower Snake Gourd —		_		Voundwort	-	41	27
	5	_		Snake Gourd		32 66	21	\mathbf{v}			
Morea		239		Snapdragon —	-	42	44 28)	Cellow Root —	-	284	190
Moon Trefoil	_		117 3	Solomon's Scal		101	67			3	7
Motherwort	-	80		Sour Sop —		35	23	Z			
Moonwort				Southernwood	-	2	2 2	Zerumbeth —	-	26	17
Mullein	-}	277	182 3	South Sea Tree - Sowbread		83	55				
	C	-//	- " "	JOHO CARL		115	771				